

Gravitational Waves from binary black holes in the extreme mass ratio regime: Self-force calculations

Carlos O. Lousto

Department of Physics and Astronomy, and Center for Gravitational Wave
Astronomy, The University of Texas at Brownsville, Brownsville, Texas 78520, USA

Abstract. We present a report on the progress made in the development of computational techniques to evaluate the gravitational radiation generated by a particle orbiting a massive black hole to second perturbative order.

PACS numbers: 04.30.-w, 04.30.Db, 04.25.Nx, 04.70.Bw

1. Introduction

It is currently believed that most of the galaxies host supermassive black holes (SMBH) in their cores [20]. Even our own galaxy host a black hole of almost 4 million solar masses [25] in its center. In this scenario, the most likely event is that stars and other compact objects in the neighborhood of the SMBH will inspiral-down and plunge toward this central massive black hole.

From the Theoretical point of view the two body problem in General Relativity remains one of the more interesting problems still unsolved. While comparable mass binary black holes are the most efficient astrophysical system regarding emission of gravitational radiation, they represent also the most challenging problem for numerical relativity. Current estimates show that during the final plunge of two nearly equal mass black holes the system radiates as much energy as in the whole previous inspiral history of the binary [4, 1, 5, 2, 3].

For the Laser Interferometric Space Antenna (LISA), the small mass ratio regime is particularly interesting since numerous sources will be 'visible' with huge signal to noise ratios [15]. This scenario is tractable using perturbation theory of black holes, but proved to be a more challenging problem than initially expected, until relatively recently, when decisive progress toward the complete solution to the problem has been made, some of which we will briefly report here.

2. The problem and its solution

The theory to compute radiation from an orbiting particle around a black hole has been developed in the 70's [23, 26, 24]. Soon after it was realized that correcting the

geodesic of the particle by its radiation formally lead to infinities. In fact, if we try a straightforward solution to the case of a particle represented by a Dirac delta in orbit around a black hole we can write the field equations of general relativity as

$$G_{\mu\nu}[h_{\alpha\beta}] = T_{\mu\nu} = m \int u_\mu u_\nu \delta[x^\alpha - x_p^\alpha(\tau)] / \sqrt{-g} d\tau \quad (1)$$

where $h_{\alpha\beta}$ represents the perturbations around the larger black hole background metric, $g_{\mu\nu}^{(0)}$ (with determinant g). $x_p^\alpha(\tau)$ is the (parametrized by the proper time of the particle, τ) coordinate location of the particle (with mass m) orbiting the large hole and $u^\mu = dx^\mu/d\tau$ is its four-velocity.

Once we solve for the perturbed metric we can correct the trajectory of the particle by requesting it to be a geodesic of the background *plus* the perturbed metric

$$\frac{d^2 x^\mu}{d\tau^2} + \Gamma_{\alpha\beta}^\mu \frac{dx^\alpha}{d\tau} \frac{dx^\beta}{d\tau} = 0. \quad (2)$$

The problem arises when trying to evaluate the connection term $\Gamma_{\alpha\beta}^\mu$ on the trajectory of the particle since it can be readily shown that the metric behaves as

$$h_{\mu\nu} \sim \frac{m H_{\mu\nu}}{\sqrt{(g_{\alpha\beta} + u_\alpha u_\beta) (x^\alpha - x_p^\alpha(\tau)) (x^\beta - x_p^\beta(\tau))}} \quad (3)$$

i.e., diverging as the inverse of the distance to the particle. So, the small perturbations assumption breaks down near the particle.

In order for us to extract any physically meaningful result this problem needs to be regularized. This proved to be a very challenging problem for over 25 years until 1997 when Mino, Sasaki and Tanaka [21] published their seminal work on the computation of the self-force a particle traveling in a curved spacetime exerts on itself. They computed this by using two alternative methods. First using matched asymptotic expansions of the perturbed background of the larger black hole by the presence of the particle and the spacetime around the small particle tidally distorted by the larger hole. By matching this two expansions in a buffer zone of radius d around the particle, with $m \ll d \ll M$, one finds that requiring the consistency of the two expansions leads to the equations of motion. The second derivation is done via a Brehme-DeWitt approach by deriving a 'conserved' rank-two symmetric tensor and integrating its divergence over the interior of the world tube surrounding the particle to obtain the equations of motion.

The self-force (defined as the correction to the background geodesic) takes then the form

$$F_{full}^\mu(x) = m \nabla^{\mu\beta\gamma} h_{\beta\gamma} = F_{direct}^\mu + F_{tail}^\mu \quad (4)$$

where F_{direct}^μ is computed from contributions that propagate *along* the past light cone and F_{tail}^μ has the contributions from *inside* the light cone, product of the scattering of perturbations due to the motion of the particle in the curved spacetime created by the larger black hole.

The self-force is then computed by taking the limit $F_{self}^\mu = F_{tail}^\mu(x \rightarrow x_p)$ where

$$F_{tail}^\mu = m u^\alpha u^\beta \int_{-\infty}^{\tau_-} \left\{ \frac{1}{2} \nabla^\mu G_{\alpha\beta\gamma'\delta'}^- - \nabla_\beta G_{\alpha\gamma'\delta'}^- u^\mu - \frac{1}{2} u^\mu u^\lambda \nabla_\lambda G_{\alpha\beta\gamma'\delta'}^- \right\} u^{\gamma'} u^{\delta'} d\tau' \quad (5)$$

and where $G_{\alpha\beta\gamma'\delta'}^-$ is the retarded Green's function. Primed indices denote evaluation at the particle's location while unprimed denotes evaluation at the nearby field point x ($u^\alpha u^\beta$ are obtained by parallel propagation.) Finally, τ_- represents the proper time of a point on the worldline that intersects the particle's past lightcone of the field point x . As a result, one finds that the divergent terms mentioned above do not affect the motion of the particle.

Independently, Quinn and Wald [22] come up with the same equation of motion (5) (usually referred to as 'MiSaTaQuWa' force), but using an axiomatic approach:

i) comparison axiom: Identifying the four-velocity u^μ and the four-acceleration at points P_1 in a manifold $(M_1, g_{\mu\nu}^{(1)})$ and P_2 in a manifold $(M_2, g_{\mu\nu}^{(2)})$ via Riemann normal coordinates one find that the difference between forces is given by

$$f_{(1)}^\mu - f_{(2)}^\mu = \lim_{r \rightarrow 0} \left\{ \left(\frac{1}{2} \nabla_{(1)}^\mu h_{\alpha\beta}^{(1)} - \nabla_{(1)\beta} h_{\alpha}^{(1)\mu} \right) - \left(\frac{1}{2} \nabla_{(2)}^\mu h_{\alpha\beta}^{(2)} - \nabla_{(2)\beta} h_{\alpha}^{(2)\mu} \right) \right\} \quad (6)$$

ii) Flat spacetime axiom: If $(M, g_{\mu\nu})$ is the Minkowski spacetime the perturbed metric is the half advanced and half retarded solution and the force vanishes

$$h_{\alpha\beta} = \frac{1}{2} [h_{\alpha\beta}^+ + h_{\alpha\beta}^-], \text{ and } f^\mu = m u^\alpha \nabla_\alpha^{(0)} u^\mu = 0. \quad (7)$$

It is worth mentioning here a newer approach [14] to the problem which decomposes the full fields into a 'regular' (which is different from the 'tail' part) and a 'singular' part. It can be proved that only the regular part contributes to the self-force, producing the same 'MiSaTaQuWa' result, and that the regular (metric) fields satisfy the vacuum equations of General Relativity (with no source terms due to the presence of the particle).

The self-force calculation thus provides a local correction to the trajectory of the particle taking into account radiative and nonradiative effects. For a global method that takes into account the radiative effects only see, for instance Ref. [16], that makes use of the energy and momentum balance of the radiation emitted to infinity and onto the black hole horizon to correct the particle trajectory over orbital cycles.

3. The problems of the solution

The proof that the equation of motion

$$m \left(\frac{d^2 x^\mu}{d\tau^2} + \Gamma_{\alpha\beta}^{(0)\mu} \frac{dx^\alpha}{d\tau} \frac{dx^\beta}{d\tau} \right) = F_{self}^\mu \quad (8)$$

leads to the correct, finite description of the particle due to its own field represent an enormous leap forward in our comprehension of the effects of gravitational radiation. Yet, expression (5) for the self-force is formal and it was not immediately ready for implementation. It took a couple of years to develop the formalism that allowed explicit computations. The key observation, arrived to simultaneously by several authors

[8, 17, 11], was to tackle the nonrotating, i.e. Schwarzschild black hole, background case first to make use of its spherical symmetry to expand the force into spherical harmonics. For each multipole $(\ell, \mathfrak{m})$, the full force is finite, and the divergent nature of the problem only reveals upon summing over $(\ell, \mathfrak{m})$.

In the mode sum regularization method the force can be decomposed as (where we performed already the sum over the modes $\mathfrak{m}$)

$$F_{self}^\mu = \lim_{x \rightarrow x_p} \sum_{\ell} \left[F_{full}^{\mu\ell}(x) - F_{direct}^{\mu\ell}(x) \right] \quad (9)$$

which upon addition and subtraction of $A^\mu L + B^\mu + C^\mu/L$ (which leads to divergences upon summing over ℓ)

$$\begin{aligned} F_{self}^\mu &= \sum_{\ell} \left[F_{full}^{\mu\ell}(x_p) - (A^\mu L + B^\mu + C^\mu/L) \right] \\ &\quad - \sum_{\ell} \left[F_{direct}^{\mu\ell}(x_p) - (A^\mu L + B^\mu + C^\mu/L) \right] \end{aligned} \quad (10)$$

where for convenience we defined $L = \ell + 1/2$. The terms in brackets now lead to well defined, convergent series. The sum over the last bracket is a remainder normally denoted as D^μ .

In a notable piece of work, the regularization parameters A^μ, B^μ, C^μ and D^μ have been computed *analytically*, first for the Schwarzschild [7] and then for the Kerr backgrounds [10].

The missing part is now the full force, $F_{full}^{\mu\ell}(x_p)$, that has to be computed by numerically integrating the (perturbed) General Relativity field equations to obtain $F_{full}^{\mu\ell}(x)$. For a nonrotating large black hole, we can just solve the wave equations for the perturbations traveling on the curved Schwarzschild background, i.e. the Zerilli and Regge-Wheeler equation for the two polarizations of the radiation. We then know how to reconstruct the metric in the Regge-Wheeler gauge [17].

We successfully followed this procedure in the case of a particle starting from rest at a finite distance from the large black hole [6]. In addition to the exact (numerical) computation of the self-force, we obtained an analytic approximation

$$F_{self}^r = F_{self}^{r\ell=0} + F_{self}^{r\ell=1} + \sum_{\ell=2}^{\infty} \left\{ -\frac{15}{16} m^2 \frac{E^2}{r^2} \left(E^2 + \frac{4M}{r} - 1 \right) \frac{1}{L^2} + \mathcal{O}(L^{-4}) \right\} \quad (11)$$

where M stands for the mass of the large hole and (m, E) are the mass and orbital energy of the particle respectively. Note also that the force will have [18] only negative, even powers of L what makes the sum quickly convergent and provides an excellent approximation if we evaluate the first few lower multipoles numerically.

After this successful computation we proceeded to compute the astrophysically more appealing scenario of a particle in circular orbit around a nonrotating black hole. In this case we encountered a fundamental difficulty: The Regge-Wheeler gauge does not lead to C^0 metric coefficients at the particle location as for headon orbits. Moreover, in general, a Dirac's delta behaviour appear in the metric coefficients at x_p . Besides, the regularization parameters have been computed in the harmonic gauge.

We have delayed until now the discussion of the important issue of the *gauge problem*. In fact, the 'MiSaTaQuWa' force (5) is computed in the harmonic gauge. Since the 'force' is not a gauge invariant concept we need to deal with the problem of gauge transformations [9]. In particular, it is difficult to compute the transformation between a singular gauge at the location of the particle, as the Regge-Wheeler gauge, and a regular and locally isotropic one as the harmonic gauge.

4. The solution to the problems of the solution

The reason we used the Regge-Wheeler gauge was its convenience for solving General Relativity's field equations, i.e. reducing them to solve the Zerilli and Regge-Wheeler wave equations. The natural way of proceeding for a computation oriented to the evaluation of the self-force is to solve the ten General Relativity field equations

$$\square \bar{h}_{\alpha\beta} + 2R^\mu{}_\alpha{}^\nu{}_\beta \bar{h}_{\mu\nu} = S_{\alpha\beta}, \quad (12)$$

where $S_{\alpha\beta}$ are the source terms due to the orbiting particle, directly in the *harmonic gauge*

$$g^{\beta\gamma} \bar{h}_{\alpha\beta;\gamma} = 0, \text{ where } \bar{h}_{\alpha\beta} = h_{\alpha\beta} - \frac{1}{2} g_{\alpha\beta} h. \quad (13)$$

The main inconvenience to overcome are the potential numerical instabilities of this formulation.

A proper use of the coordinate conditions (13) that eliminates all the first time derivatives leading to instabilities in equations (12) is possible (Barack and Lousto, in preparation). We can then use the algorithm of Ref. [19] to integrate the wave equations with the Dirac's delta source terms appearing in (12).

The advantages of this procedure are apparent: No gauge problem, the computed force will be in the harmonic gauge; so are the regularization parameters. We have to deal with C^0 quantities (metric coefficients) as opposed to discontinuous waveforms or divergent metric coefficients, and will only need first order derivatives of them to compute the self-force. The procedure is in principle generalizable to Kerr backgrounds (although the actual implementation will not be a straightforward one). Finally, the non-radiative modes $\ell = 0, 1$ can be solved in practically an analytic way in the harmonic gauge [13].

5. more problems seeking for solutions

As we mentioned earlier the self-force is not a gauge invariant concept. A way to recover a physical interpretation of the corrected trajectory of the particle is to use it to compute the corrections to the energy and momentum radiated at infinity from the binary system. In order to do that we have to deal with second order perturbations of the large black hole background. Including this metric perturbations is needed to consistently take into account all $\mathcal{O}(m/M)^2$ terms (the other coming from the corrected trajectory of

the particle.) To second perturbative order, products of first order perturbations act as sources [12], as a result, we could have singular products at the location of the particle (we only probed regularization of the force, not of each component of the first order metric and its derivatives).

Rosenthal and Ori have recently worked out a scheme to extract the leading singular piece of the (quadratic) source term using the retarded field and a choice of the local gauge to regularize the problem. This promising approach may solve the last standing theoretical problem, but we know by now that new, sometimes unexpected, problems may still arise in our way to provide the complete solution.

Acknowledgments

The author thanks L.Barack for reading the original manuscript. We also gratefully acknowledge the support of the NASA Center for Gravitational Wave Astronomy at The University of Texas at Brownsville (NAG5-13396) and NSF for grants PHY-0140326 and PHY-0354867.

- [1] J. Baker, B. Brügmann, Manuela Campanelli, C. O. Lousto, and R. Takahashi, *Plunge waveforms from inspiralling binary black holes*, Phys. Rev. Lett. **87** (2001), 121103.
- [2] J. Baker, M. Campanelli, C. O. Lousto, and R. Takahashi, *Modeling gravitational radiation from coalescing binary black holes*, Phys. Rev. D **65** (2002), 124012.
- [3] ———, *Coalescence remnant of spinning binary black holes*, Phys. Rev. **D69** (2004), 027505.
- [4] John Baker, Bernd Brügmann, Manuela Campanelli, and Carlos O. Lousto, *Gravitational waves from black hole collisions via an eclectic approach*, Class. Quantum Grav. **17** (2000), L149–L156.
- [5] John Baker, Manuela Campanelli, and Carlos O. Lousto, *The Lazarus project: A pragmatic approach to binary black hole evolutions*, Phys. Rev. D **65** (2002), 044001.
- [6] Leor Barack and Carlos O. Lousto, *Computing the gravitational self-force on a compact object plunging into a schwarzschild black hole*, Phys. Rev. **D66** (2002), 061502.
- [7] Leor Barack, Yasushi Mino, Hiroyuki Nakano, Amos Ori, and Misao Sasaki, *Calculating the gravitational self force in schwarzschild spacetime*, Phys. Rev. Lett. **88** (2002), 091101.
- [8] Leor Barack and Amos Ori, *Mode sum regularization approach for the self force in black hole spacetime*, Phys. Rev. D **61** (2000), 061502.
- [9] ———, *Gravitational self force and gauge transformations*, Phys. Rev. D **64** (2001), 124003.
- [10] ———, *Gravitational self-force on a particle orbiting a kerr black hole*, Phys. Rev. Lett. **90** (2003), 111101.
- [11] Lior M. Burko, *Self force on static charges in schwarzschild spacetime*, Class. Quant. Grav. **17** (2000), 227–250.
- [12] M. Campanelli and C. O. Lousto, *Second order gauge invariant gravitational perturbations of a Kerr black hole*, Phys. Rev. D **59** (1999), 124022.
- [13] Steven Detweiler and Eric Poisson, *Low multipole contributions to the gravitational self- force*, Phys. Rev. **D69** (2004), 084019.
- [14] Steven Detweiler and Bernard F. Whiting, *Self-force via a green’s function decomposition*, Phys. Rev. **D67** (2003), 024025.
- [15] Jonathan R. Gair et al., *Event rate estimates for lisa extreme mass ratio capture sources*, Class. Quant. Grav. **21** (2004), S1595–S1606.
- [16] Scott A. Hughes, *Evolution of circular, non-equatorial orbits of Kerr black holes due to gravitational-wave emission. II: Inspiral trajectories and gravitational waveforms*, Phys. Rev. D **64** (2001), 064004.

- [17] Carlos O. Lousto, *Pragmatic Approach to Gravitational Radiation Reaction in Binary Black Holes*, Phys. Rev. Lett. **84** (2000), 5251.
- [18] ———, *Towards the solution of the relativistic gravitational radiation reaction problem for binary black holes*, Class. Quantum Grav. **18** (2001), 3989.
- [19] Carlos O. Lousto and Richard H. Price, *Understanding initial data for black hole collisions*, Phys. Rev. D **56** (1997), 6439–6457.
- [20] David Merritt and Milos Milosavljevic, *Massive black hole binary evolution*, (2004).
- [21] Yasushi Mino, Misao Sasaki, and Takahiro Tanaka, *Gravitational radiation reaction to a particle motion*, Phys. Rev. D **55** (1997), 3457–3476.
- [22] Theodore C. Quinn and Robert M. Wald, *An axiomatic approach to electromagnetic and gravitational radiation reaction of particles in curved spacetime*, Phys. Rev. D **56** (1997), 3381–3394.
- [23] T. Regge and J. Wheeler, *Stability of a Schwarzschild singularity*, Phys. Rev. **108** (1957), no. 4, 1063–1069.
- [24] S. A. Teukolsky, *Perturbations of a rotating black hole. I. fundamental equations for gravitational, electromagnetic, and neutrino-field perturbations*, Astrophys. J. **185** (1973), 635–647.
- [25] Nevin N. Weinberg, Milos Milosavljevic, and Andrea M. Ghez, *Stellar dynamics at the galactic center with a thirty meter telescope*, (2004).
- [26] F. J. Zerilli, *Effective potential for even-parity Regge-Wheeler gravitational perturbation equations*, Phys. Rev. Lett. **24** (1970), no. 13, 737–738.